

SOMAÍ THC800 Indica GS

Consumer Medicine Information (CMI) summary

The [full CMI](#) on the next page has more details. If you are worried about using this medicine, speak to your doctor or pharmacist.

▼ This medicine is new and is not yet approved on the Australian Register of Therapeutic Goods. Please report side effects. See the [full CMI](#) for further details.

WARNING: SOMAÍ THC800 Indica GS is an unapproved medicine that contains controlled substances. Possession without authority is illegal.

1. WHY AM I USING SOMAÍ THC800 INDICA GS?

SOMAÍ THC800 Indica GS is an inhalation cartridge that contains the active ingredient tetrahydrocannabinol (THC). Each 1mL of **SOMAÍ THC800 Indica GS** contains 800mg THC.

Your doctor can indicate the desired symptom or condition for which this medication has been prescribed to you. For more information, see Section [1. Why am I using SOMAÍ THC800 Indica GS?](#) in the full CMI.

2. WHAT SHOULD I KNOW BEFORE I USE SOMAÍ THC800 INDICA GS?

Do not use if you have ever had an allergic reaction to THC, CBD, any other known cannabinoid or any of the ingredients listed at the end of the CMI.

Talk to your doctor if you have any other medical conditions, take any other medicines, or are pregnant or plan to become pregnant or are breastfeeding. For more information, see Section [2. What should I know before I use SOMAÍ THC800 Indica GS?](#) in the full CMI.

3. WHAT IF I AM TAKING OTHER MEDICINES?

Some medicines may interfere with **SOMAÍ THC800 Indica GS** and affect how it works.

A list of these medicines is in Section [3. What if I am taking other medicines?](#) in the full CMI.

4. HOW DO I USE SOMAÍ THC800 INDICA GS?

- There is no scientifically defined dose of medical cannabis for any specific medical condition. Use **SOMAÍ THC800 Indica GS** exactly as prescribed by your doctor.
- Your doctor will assess you and give you a dosing plan to follow.

More instructions can be found in Section [4. How do I use SOMAÍ THC800 Indica GS?](#) in the full CMI.

5. WHAT SHOULD I KNOW WHILE USING SOMAÍ THC800 INDICA GS?

Things you should do	<ul style="list-style-type: none">• Remind any doctor, dentist or pharmacist you visit that you are using SOMAÍ THC800 Indica GS.• Follow the dosing instructions provided by your doctor, because THC can produce dependence, presenting the potential for abuse.• Check with the country or countries that you intend to visit or crossing for information on whether you may take your medicine, because SOMAÍ THC800 Indica GS contains controlled substances.
Things you should not do	<ul style="list-style-type: none">• Do not stop using this medicine suddenly.• Do not give this medicine to anyone else, even if they have the same condition as you.
Driving or using machines	<ul style="list-style-type: none">• Do not drive or operate machinery while being treated with SOMAÍ THC800 Indica GS.• Be aware that in most Australian states and territories, driving with detectable levels of THC in your body is illegal, even though it may not impair your driving.
Drinking alcohol	<ul style="list-style-type: none">• Tell your doctor if you drink alcohol.• You should avoid drinking alcohol while using SOMAÍ THC800 Indica GS.
Looking after your medicine	<ul style="list-style-type: none">• Store below 25°C. Store in the original package in order to protect from light and moisture.• Keep it where children cannot reach it.

For more information, see Section [5. What should I know while using SOMAÍ THC800 Indica GS?](#) in the full CMI.

6. ARE THERE ANY SIDE EFFECTS?

Call your doctor straight away, or go straight to the nearest hospital if you notice any of these serious side effects, such as: signs of an allergic reaction, paranoia, hallucinations, fast or irregular heartbeat, low blood pressure if associated with lightheadedness, dizziness or fainting, seizures, depression or thoughts of suicide. Less serious side effects can occur and you should speak to your doctor if any of them worry you.

For more information, including what to do if you have any side effects, see Section [6. Are there any side effects?](#) in the full CMI.

SOMAÍ THC800 Indica GS

Active ingredient(s): *Tetrahydrocannabinol (THC)*

Consumer Medicine Information (CMI)

This leaflet provides important information about using **SOMAÍ THC800 Indica GS**. You should also speak to your doctor or pharmacist if you would like further information or if you have any concerns or questions about using **SOMAÍ THC800 Indica GS**.

Where to find information in this leaflet:

1. Why am I using **SOMAÍ THC800 Indica GS**?
2. What should I know before I use **SOMAÍ THC800 Indica GS**?
3. What if I am taking other medicines?
4. How do I use **SOMAÍ THC800 Indica GS**?
5. What should I know while using **SOMAÍ THC800 Indica GS**?
6. Are there any side effects?
7. Product details

1. WHY AM I USING SOMAÍ THC800 INDICA GS?

SOMAÍ THC800 Indica GS contains the active ingredient **tetrahydrocannabinol (THC)**. Each 1mL of **SOMAÍ THC800 Indica GS** contains 800mg THC.

SOMAÍ THC800 Indica GS contains ingredients called cannabinoids, which have been extracted from cannabis plants. Cannabinoids, such as THC and CBD, interact with the body's endocannabinoid system to produce various effects. The therapeutic benefits of medicinal cannabis may vary depending on a patient's biology and the product selected. **SOMAÍ THC800 Indica GS** is high in THC.

SOMAÍ THC800 Indica GS is not approved for treatment of any specific conditions or symptoms. Your doctor can indicate the desired symptom or condition for which this medication has been prescribed to you.

To date, the Therapeutic Goods Administration (TGA) has approved medicinal cannabis SAS applications for indications including, but not limited to, the following:

- chemotherapy-induced nausea and vomiting
- refractory paediatric epilepsy
- symptom relief in palliative care, including non-cancer related palliation
- cancer pain
- neuropathic pain, including that related to multiple sclerosis
- spasticity from neurological conditions, such as multiple sclerosis
- anorexia and wasting associated with chronic illness
- Parkinson's Disease
- mental health disorders

Your doctor may also prescribe this medication for conditions or symptoms which have not been mentioned.

Ask your doctor if you have any questions about why **SOMAÍ THC800 Indica GS** has been prescribed for you.

2. WHAT SHOULD I KNOW BEFORE I USE SOMAÍ THC800 INDICA GS?

Warnings

Do not use **SOMAÍ THC800 Indica GS** if:

- you are allergic to THC, CBD, any other known cannabinoid, or any of the ingredients listed at the end of this leaflet. Always check the ingredients to make sure you can use this medicine
- you or a close family member directly related to you has any mental health conditions including psychosis, schizophrenia, or other significant psychiatric disorder
- you have severe and unstable cardio-pulmonary disease (angina, peripheral vascular disease, cerebrovascular disease and arrhythmias) or risk factors for heart disease
- you have severe kidney or liver conditions
- you are pregnant or breast-feeding, think you may be pregnant or are planning to become pregnant
- you have an underlying pulmonary disease such as emphysema, chronic bronchitis, pulmonary infections

Check with your doctor if you:

- have mild or moderate kidney or liver conditions because your doctor may consider that you need to be monitored while using this medicine
- have a history of drug dependence, including nicotine and heavy users of alcohol
- have a history of epilepsy or recurrent seizures due to the THC content of this medicine
- take any medicines for any other condition. See additional information under *Section 3. What if I am taking other medicines?*
- are an elderly patient (more than 65 years of age) because you may be more sensitive to the neurological, psychoactive, and postural hypotensive effects of cannabinoids
- are a younger patient (under 18 years of age) due to the potentially adverse effects of cannabis on the developing brain
- suffer from dyspnea or a condition that makes inhalation or pulmonary absorption difficult because in these cases this route of administration should not be used

The use of **SOMAÍ THC800 Indica GS** may involve additional health risks beyond those described above, some of which are not known or fully understood. There may be additional contraindications/precautions that are unknown due to limited medical and scientific information.

During treatment, you may be at risk of developing certain side effects. It is important you understand these risks and how to monitor for them. See additional information under Section 6. *Are there any side effects?*

Pregnancy and breastfeeding

Check with your doctor if you are pregnant or intend to become pregnant.

Talk to your doctor if you are breastfeeding or intend to breastfeed.

If you are pregnant or breast-feeding, think you may be pregnant or are planning to become pregnant you cannot take this medicine.

Fertility

Cannabis preparations should not be used by anyone planning to have a child. Cannabinoids may impact fertility (whether male or female) and can affect the gestational development of the fetus.

Check with your doctor if you are planning to have a baby.

It is advisable to always use a highly effective contraceptive method. Women using hormonal contraceptives should add a barrier non-hormonal contraceptive method during treatment with **SOMAÍ THC800 Indica GS**.

● 3. WHAT IF I AM TAKING OTHER MEDICINES?

Tell your doctor or pharmacist if you are taking any other medicines, including any medicines, vitamins or supplements that you buy without a prescription from your pharmacy, supermarket or health food shop.

Some medicines may interact with SOMAÍ THC800 Indica GS and affect how it works, increasing or reducing its effects. SOMAÍ THC800 Indica GS may also interact with some medicines and affect how they work.

There is limited evidence about how medicinal cannabis reacts with other approved medications. Most interactions are associated with drugs that slow down the nervous system, causing drowsiness, such as sedatives, benzodiazepines and hypnotics. The list below does not include all possible drug interactions. The doctor or pharmacist should evaluate all possible drug interactions on a case-by-case basis.

- The association with antispastic medications, used in muscle-related diseases, can reduce muscle strength and increase the risk of falls, so precautions must be taken
- Medicines used for sleep problems (sleeping pills or sedatives/hypnotics such as: benzodiazepines, zopiclone, zolpidem)
- Medications used for pain (opioids, such as oxycodone, hydrocodone, codeine and morphine)
- Medicines used to treat epilepsy, seizures or neuropathic pain (for example, carbamazepine, phenytoin, primidone, phenobarbital, clobazam)
- Medicines used to treat bacterial (e.g. antibiotics such as rifampicin, rifabutin, clarithromycin and erythromycin) or fungal (e.g. antifungals such as

itraconazole, fluconazole, ketoconazole and miconazole, amphotericin B) infections

- Medicines used to thin the blood (anticoagulants such as warfarin, apixaban, rivaroxaban)
- Medicines used for heart disease or hypertension (such as amiodarone, diltiazem and verapamil)
- Medicines used to control HIV/AIDS infection (HIV protease inhibitors) in particular ritonavir
- Medicines used to treat diabetes (troglitazone)
- St John's Wort
- Anti-depressants (fluoxetine, fluvoxamine)
- Stomach acid inhibitors (omeprazole)
- Medicines that are strongly bound to proteins (e.g. cyclosporine)
- Any drug or complementary medicine metabolised by cytochrome P450

If you stop taking any of these medications, it may cause an increase in the bioavailability of THC, and there may be a need to adjust the dose titration to be administered.

Check with your doctor or pharmacist if you are not sure about what medicines, vitamins or supplements you are taking and if these affect SOMAÍ THC800 Indica GS.

If at any time you suspect that a combination of medicines is working in an unexpected way, or has any unwanted or unpleasant effects, consult your doctor immediately.

● 4. HOW DO I USE SOMAÍ THC800 INDICA GS?

How much to take

Treatment must be initiated and supervised by a doctor.

There is no scientifically defined dose of medical cannabis for any specific medical condition. There are no set dosage schedules for medical cannabis; dose titration is necessary and requires an individualized approach to adjust the ideal dosage needed for each patient and the final dosage depends on the person and the condition being treated.

Your doctor will work with you to decide the dose of **SOMAÍ THC800 Indica GS** that best suits you and your symptoms and minimizes any side effects. It is highly recommended to begin at the lowest possible dose and increase slowly over time.

Follow the instructions provided by your doctor. Check with your doctor or pharmacist if you are not sure.

When to take SOMAÍ THC800 Indica GS

Once your ideal dosage of **SOMAÍ THC800 Indica GS** is found for your condition, continue to take that dose as needed.

Follow the instructions provided and use this medicine until your doctor tells you to stop.

How to use SOMAÍ THC800 Indica GS

SOMAÍ THC800 Indica GS is for inhalation use.

SOMAÍ THC800 Indica GS cartridge is designed for inhalation with a compatible vaporisation device and should not be used with any other devices.

Be aware that the temperature setting with vaporisers will impact the effects of this medicine, as each cannabinoid has a different boiling point.

Use the vaporiser in line with the manufacturer's instructions or seek advice from your doctor or pharmacist.

If you forget to use SOMAÍ THC800 Indica GS

SOMAÍ THC800 Indica GS should be used as directed by your doctor.

If you miss your dose at the usual time or if it is almost time for your next dose, skip the dose you missed and take your next dose when you are meant to.

Do not take a double dose to make up for the dose you missed.

If you use too much SOMAÍ THC800 Indica GS

If you think that you have used too much **SOMAÍ THC800 Indica GS**, you may need urgent medical attention.

You should immediately:

- phone the Poisons Information Centre (**by calling 13 11 26**), or
- contact your doctor, or
- go to the Emergency Department at your nearest hospital.

You should do this even if there are no signs of discomfort or poisoning.

5. WHAT SHOULD I KNOW WHILE USING SOMAÍ THC800 INDICA GS?

Things you should do

Talk to your doctor if your condition worsens or does not improve.

- THC can produce physical and psychological dependence, presenting the potential for abuse, although in the case of medical treatment with low doses of cannabis this is very unlikely to occur. You should follow the dosing instructions provided by your doctor.

Call your doctor straight away if you:

- feel unwell or experience any side effects that worry you while using your **SOMAÍ THC800 Indica GS**. See additional information under Section 6. *Are there any side effects?*
- stop taking **SOMAÍ THC800 Indica GS** suddenly

Remind any doctor, dentist or pharmacist you visit that you are using **SOMAÍ THC800 Indica GS**.

Things you should not do

- Do not stop using this medicine suddenly. If this happens, the following symptoms may occur, especially within the first 7 days: anger or aggression, irritability, anxiety, nightmares, strange dreams, insomnia, sleep disturbances, desire to consume, headache, restlessness, decreased appetite or weight loss,

depressed mood, chills, stomach pain, tremors and sweating.

- Do not give **SOMAÍ THC800 Indica GS** to anyone else, even if they have the same condition as you.

Travelling overseas with SOMAÍ THC800 Indica GS

- As **SOMAÍ THC800 Indica GS** contains controlled substances, the legality of its use and transport may differ between countries. As such, you should check if the country or countries that you intend to visit or crossing for information on whether you may take your medicine.

Driving or using machines

Do not drive or operate machinery while being treated with SOMAÍ THC800 Indica GS.

THC can cause drowsiness or dizziness and affect cognitive abilities. Such effects may impact concentration, coordination, delay reaction time and thus impair motor skills. These effects can be further increased when taking other medications or drinking alcohol.

Talk to your doctor about driving, using machines, or when undertaking other activities that require balance such as cycling or other sports, because this medicine may affect your ability to do so safely.

Measurable concentrations of THC (the main psychoactive substance in cannabis) can be detected in urine many days after the last dose of **SOMAÍ THC800 Indica GS**. It may take up to 5 days for 80 to 90% of the dose to be excreted. Be aware that in most Australian states and territories, driving with detectable levels of THC in your body is illegal, even though it may not impair your driving.

Drinking alcohol

Tell your doctor if you drink alcohol.

You should avoid drinking alcohol while using **SOMAÍ THC800 Indica GS**.

Alcohol can increase the effects of **SOMAÍ THC800 Indica GS** that impair cognitive abilities, reaction time, coordination and motor skills. Cognitive impairment may be greatly increased when **SOMAÍ THC800 Indica GS** is consumed along with alcohol or other drugs which affect the activity of the nervous system (e.g. opioids, sleeping pills, other psychoactive drugs).

Looking after your medicine

Follow the instructions in the pack on how to take care of your medicine properly.

Store below 25°C. Store in the original package in order to protect from light and moisture.

Keep it where children cannot reach it.

When to discard your medicine

Do not use this medicine if you notice any visible signs of deterioration.

Getting rid of any unwanted medicine

If you no longer need to use this medicine or it is out of date, take it to any pharmacy for safe disposal.

Do not use this medicine after the expiry date, which is stated on the pack. The expiry date refers to the last day of that month.

6. ARE THERE ANY SIDE EFFECTS?

All medicines can have side effects. If you do experience any side effects, most of them are minor and temporary. However, some side effects may need medical attention.

See the information below and, if you need to, ask your doctor or pharmacist if you have any further questions about side effects.

Information on the side effects associated with medical cannabis use is limited. Patients generally tolerate it well. Often a low dosage provides sufficient relief so that side effects rarely occur. When these happen, they are usually the result of a high dosage or combined use with a substance, such as alcohol, which intensifies its effects.

The toxic and pharmacological effects of cannabinoids may overlap due to the small therapeutic window of THC, which is mainly responsible for possible side effects.

Medicinal cannabis side effects are commonly dose-dependent, so it's important to follow your doctor's dosing recommendations.

Long-term use may decrease one or more of the effects of cannabis (tolerance).

Less serious side effects

Less serious side effects	What to do
<ul style="list-style-type: none">● Dizziness● Feeling sick or nauseous● Increased heart rate or palpitations● Changes in appetite● Dry mouth● Muscle weakness● Mood changes● General feeling of discomfort● Blurred vision● Anxiety● Confusion● Problems with memory, concentration, perception of reality, sense of time, ability to think and make decisions, reaction time and coordination● Problems with balance or falls	<p>Speak to your doctor if you have any of these less serious side effects and they worry you.</p>

Serious side effects

Serious side effects	What to do
<ul style="list-style-type: none">● Signs of an allergic reaction such as swelling of the face, lips, tongue or other body part, severe fainting or dizziness, or rash/itching of the skin	<p>Call your doctor straight away, or go straight to the</p>

<ul style="list-style-type: none">● Paranoia (feeling suspicious, angry or betrayed by other people)● Hallucinations (hearing voices or seeing things that aren't there)● Fast or irregular heartbeat● Low blood pressure if associated with lightheadedness, dizziness or fainting● Seizures (if you have no history of epilepsy)● Depression or thoughts of suicide	<p>Emergency Department at your nearest hospital if you notice any of these serious side effects.</p>
--	--

Tell your doctor or pharmacist if you notice anything else that may be making you feel unwell.

Other side effects not listed here may occur in some people.

Reporting side effects

After you have received medical advice for any side effects you experience, you can report side effects to the Therapeutic Goods Administration online at www.tga.gov.au/reporting-problems. By reporting side effects, you can help provide more information on the safety of this medicine.

Always make sure you speak to your doctor or pharmacist before you decide to stop taking any of your medicines.

7. PRODUCT DETAILS

This medicine is only available with a doctor's prescription.

What SOMAÍ THC800 Indica GS contains

Active ingredients (main ingredient)	<ul style="list-style-type: none">● THC: 800mg/mL
Other ingredients (inactive ingredients)	<ul style="list-style-type: none">● Hemp derived terpenes

Do not take this medicine if you are allergic to any of these ingredients.

What SOMAÍ THC800 Indica GS looks like

SOMAÍ THC800 Indica GS is a yellow to amber viscous liquid or solid at room temperature for inhalation use supplied in a 1 mL glass cartridge, inside a plastic tube with a closure, and leaflet and is available in packs of 1 unit.

Who distributes SOMAÍ THC800 Indica GS

Somaí Pharmaceuticals Pty Ltd
contact@somaipharma.com.au

This leaflet was prepared in October 2024.

SPK0174/00